

**Probability Generating Functions** are a useful tool with reference to *discrete* probability distributions, that is probabilities  $P(x)$  that are a function of *the positive integers*  $x = 0, 1, 2, 3, 4, 5, \dots$  i.e. *Uniform, Geometric, Binomial and Poisson Distributions*, but *not* a Normal distribution of *continuous* random variables (i.e. where  $x$  can take any real value).

**Definition:**  $G_x(z)$  is the Probability Generating Function (PGF) associated with discrete random variable  $x$ .  $p(x)$  is the probability of  $x$  occurring.  $z$  is a real number and  $G_x(z)$  is the expectation of the quantity  $z^x$ .

$$G_x(z) = E[z^x] = \sum_{x=0}^{\infty} p(x)z^x$$

If we can construct  $G_x(z)$  for a particular probability distribution, then its algebraic properties enable us to *efficiently compute useful properties of the distribution* (such as the *mean*, *variance* and the distribution of *combinations* of random variables). It works very much like a *Moment Generating Function* (MGF), but is *restricted to discrete random variables*.

$$G_x(1) = \sum_{x=0}^{\infty} p(x) = 1$$

by *definition* of the probability distribution  $p(x)$

### Mean

$$\frac{\partial G_x}{\partial z} \Big|_{z=1} = \sum_{x=0}^{\infty} xp(x)z^{x-1} \Big|_{z=1} = \sum_{x=0}^{\infty} xp(x) = E[x]$$

$$\therefore \mu = E[x] = G'_x(1)$$

### Variance

$$\frac{\partial^2 G_x}{\partial z^2} \Big|_{z=1} = \sum_{x=0}^{\infty} x(x-1)p(x)z^{x-2} \Big|_{z=1} = \sum_{x=0}^{\infty} x^2 p(x) - \sum_{x=0}^{\infty} xp(x) = V[x] + \mu^2 - \mu$$

$$\therefore \sigma^2 = V[x] = G''_x(1) - \mu^2 + \mu$$

### Example: Uniform distribution

$$p(x) = \frac{1}{N}$$

$$x = 1 \dots N$$

$$G_x(z) = \frac{1}{N}z^1 + \frac{1}{N}z^2 + \frac{1}{N}z^3 + \dots + \frac{1}{N}z^N$$

$$G'_x(z) = \frac{\partial G_x}{\partial z} = \frac{1}{N}z^0 + \frac{2}{N}z^1 + \frac{3}{N}z^2 + \dots + \frac{N}{N}z^{N-1}$$

$$G'_x(1) = \frac{1}{N} + \frac{2}{N} + \dots + \frac{N}{N} = \frac{1}{2}N\left(\frac{1}{N} + \frac{N}{N}\right) = \frac{1}{2}(N+1) \quad \text{Sum of arithmetic progression}$$

$$\therefore \mu = \frac{1}{2}(N+1) \quad \text{Mean}$$

$$G''_x(z) = \frac{\partial^2 G_x}{\partial z^2} = \frac{2}{N}z^0 + \frac{6}{N}z^1 + \dots + \frac{N(N-1)}{N}z^{N-2}$$

$$G''_x(1) = \frac{2}{N} + \frac{6}{N} + \dots + \frac{N(N-1)}{N} = \frac{1}{N} \sum_{n=1}^{N-1} n(n+1) = \frac{1}{N} \sum_{n=1}^{N-1} (n^2 + n)$$

$$\sum_{n=1}^N n = \frac{1}{2}N(N+1)$$

Quote these summation formulae

$$\sum_{n=1}^N n^2 = \frac{1}{6}N(2N+1)(N+1)$$

$$\frac{1}{N} \sum_{n=1}^{N-1} (n^2 + n) = \frac{1}{N} \left( \frac{1}{6}N(2N+1)(N+1) + \frac{1}{2}N(N+1) - N^2 - N \right)$$

$$\frac{1}{N} \sum_{n=1}^{N-1} (n^2 + n) = \frac{1}{6}(2N+1)(N+1) + \frac{1}{2}(N+1) - (N+1)$$

$$\frac{1}{N} \sum_{n=1}^{N-1} (n^2 + n) = \frac{1}{12}(N+1)\{4N+2+6-12\}$$

$$\frac{1}{N} \sum_{n=1}^{N-1} (n^2 + n) = \frac{1}{12}(N+1)(4N-4)$$

$$\therefore \sigma^2 = \frac{1}{12}(N+1)(4N-4) + \frac{1}{2}(N+1) - \frac{1}{4}(N+1)^2$$

$$\sigma^2 = \frac{1}{12}(N+1)(4N-4+6-3N-3)$$

$$\sigma^2 = \frac{1}{12}(N+1)(N-1) \quad \text{Variance}$$

$$\mu = E[x] = G'_x(1)$$

$$\sigma^2 = V[x] = G''_x(1) - \mu^2 + \mu$$

### Example: non uniform distribution

$$x=1,2,3$$

$$p(1)=0.5$$

$$p(2)=0.2$$

$$p(3)=0.3$$

$$G_x(z) = 0.5z + 0.2z^2 + 0.3z^3$$

$$G'(z) = 0.5 + 0.4z + 0.9z^2$$

$$G''(z) = 0.4 + 1.8z$$

$$\mu = G'_x(1) = 0.5 + 0.4 + 0.9 = 1.8$$

$$\sigma = G''_x(1) - \mu^2 + \mu = 0.4 + 1.8 - 1.8^2 + 1.8 = 0.76$$

### Example: Geometric Distribution

$$x=1, \dots, \infty$$

$$P(x) = (1-p)^{x-1} p$$

$$G_x(z) = pz + (1-p)pz^2 + (1-p)^2 pz^3 + \dots + (1-p)^{N-1} pz^N + \dots$$

Geometric progression with first term  $pz$  and common ratio  $(1-p)z$

$$G_x(z) = \lim_{N \rightarrow \infty} \left( \frac{pz(1-(1-p)^N z^N)}{1-(1-p)z} \right)$$

$$G_x(z) = \frac{pz}{1-(1-p)z}, \quad s.t. \quad |(1-p)z| < 1 \quad \text{Unless } p = 1, \text{ this will always be true when } z = 1 \text{ since } p \text{ is in the range } [0,1]$$

$$G'_x(z) = \frac{(1-(1-p)z)p - pz(-(1-p))}{(1-(1-p)z)^2} = \frac{p - pz + p^2 z + pz - p^2 z}{(1-(1-p)z)^2} = \frac{p}{(1-(1-p)z)^2}$$

$$G''_x(z) = \frac{-2p}{(1-(1-p)z)^3} (-1-p) = \frac{2p(1-p)}{(1-(1-p)z)^3}$$

$$G'_x(1) = \frac{p}{p^2} = \frac{1}{p}$$

$$G''_x(1) = \frac{2p(1-p)}{p^3} = \frac{2(1-p)}{p^2}$$

$$\therefore \mu = \frac{1}{p}$$

$$\therefore \sigma^2 = \frac{2(1-p)}{p^2} - \frac{1}{p^2} + \frac{1}{p} = \frac{2-2p-1+p}{p^2} = \frac{1-p}{p^2}$$

$$\mu = E[x] = G'_x(1)$$

$$\sigma^2 = V[x] = G''_x(1) - \mu^2 + \mu$$

### Example: Binomial Distribution

$$x=0,1,2,\dots,N$$

$$p(x|N,p) = \frac{N!}{(N-x)!x!} p^x (1-p)^{N-x}$$

$$G_x(z) = \sum_{x=0}^N p(x|N,p) z^x$$

$$G_x(z) = \frac{N!}{(N-0)!0!} (pz)^0 (1-p)^N + \frac{N!}{(N-1)!1!} (pz)^1 (1-p)^{N-1} + \dots$$

$$\dots + \frac{N!}{(N-N)!N!} (pz)^N (1-p)^{N-N}$$

$$G_x(z) = (pz + 1-p)^N$$

$$G'_x(z) = Np(pz + 1-p)^{N-1}$$

$$G''_x(z) = N(N-1)p^2(pz + 1-p)^{N-2}$$

$$\mu = G'_x(1) = Np$$

$$\sigma = G''_x(1) - \mu^2 + \mu = N(N-1)p^2 - N^2 p^2 + Np = -Np^2 + Np = Np(1-p)$$

### Example: Poisson Distribution

$$x=0,1,2,\dots$$

$$p(x|\lambda) = \frac{\lambda^x}{x!} e^{-\lambda}$$

$$G_x(z) = \sum_{x=0}^N p(x|\lambda) z^x$$

$$G_x(z) = e^{-\lambda} \left( \frac{(\lambda z)^0}{0!} + \frac{(\lambda z)^1}{1!} + \dots + \frac{(\lambda z)^x}{x!} + \dots \right)$$

$$G_x(z) = e^{-\lambda} e^{\lambda z} = e^{\lambda(z-1)}$$

$$G'_x(z) = \lambda e^{\lambda(z-1)}$$

$$G''_x(z) = \lambda^2 e^{\lambda(z-1)}$$

$$\mu = G'_x(1) = \lambda$$

$$\sigma = G''_x(1) - \mu^2 + \mu = \lambda^2 - \lambda^2 + \lambda = \lambda$$

$$\sigma = \lambda$$

## Combining discrete random variables

$$y = \sum_{i=1}^n a_i x_i \quad \text{Consider a weighted sum of random variables } x_1, x_2, \dots, x_n$$

$$\therefore G_y(z) = E\left[z^y\right] = E\left[z^{\sum_{i=1}^n a_i x_i}\right] = E\left[z^{a_1 x_1} z^{a_2 x_2} \dots z^{a_n x_n}\right]$$

If two random variables have zero covariance, this means there is no correlation between them. In this case we say each random variable is *independent*.

$$\text{cov}[x, y] = E[xy] - E[x]E[y]$$

$$\text{cov}[x, y] = 0 \Rightarrow E[xy] = E[x]E[y]$$

Therefore if  $n$  random variables  $x_1, x_2, \dots, x_n$  are all *independent*

$$G_y(z) = E\left[z^{a_1 x_1} z^{a_2 x_2} \dots z^{a_n x_n}\right] = E\left[z^{a_1 x_1}\right] E\left[z^{a_2 x_2}\right] \dots E\left[z^{a_n x_n}\right]$$

$$\therefore G_y(z) = G_{x_1}(z^{a_1}) \times G_{x_2}(z^{a_2}) \times \dots \times G_{x_n}(z^{a_n})$$

So if random variables  $x, y$  are *independent*

$$G_{x+y}(z) = G_x(z) \times G_y(z)$$

### Example: Sum of two Poisson distributed variables

$$x, y = 0, 1, 2, \dots$$

$$G_x(z) = e^{\lambda_1(z-1)}$$

$$G_y(z) = e^{\lambda_2(z-1)}$$

$$G_{x+y}(z) = G_x(z) \times G_y(z)$$

$$G_{x+y}(z) = e^{(\lambda_1 + \lambda_2)(z-1)}$$

$$\mu = G_x(1) = \lambda_1 + \lambda_2$$

$$\sigma^2 = G_x''(1) - \mu^2 + \mu = (\lambda_1 + \lambda_2)^2 - (\lambda_1 + \lambda_2)^2 + \lambda_1 + \lambda_2$$

$$\sigma = \sqrt{\lambda_1 + \lambda_2}$$

$$X \sim \text{Po}(\lambda_1), \quad Y \sim \text{Po}(\lambda_2)$$

$$X + Y \sim \text{Po}(\lambda_1 + \lambda_2)$$

Which proves the result:

### Example: Sum of two six sided dice being rolled

Outcomes for each dice are 1, 2, 3, 4, 5, 6. Probabilities for each one are  $1/6$ . Let  $x$  be the outcome of dice 1 and  $y$  be the outcome of dice 2. Assume both are independent.

$$G_x(z) = \frac{1}{6}z^1 + \frac{1}{6}z^2 + \frac{1}{6}z^3 + \frac{1}{6}z^4 + \frac{1}{6}z^5 + \frac{1}{6}z^6$$

$$G_y(z) = \frac{1}{6}z^1 + \frac{1}{6}z^2 + \frac{1}{6}z^3 + \frac{1}{6}z^4 + \frac{1}{6}z^5 + \frac{1}{6}z^6$$

The PGF for the sum of  $x$  and  $y$  is therefore:

$$G_{x+y}(z) = G_x(z) \times G_y(z)$$

$$G_{x+y}(z) = \frac{1}{36}(z^1 + z^2 + z^3 + z^4 + z^5 + z^6)^2$$

$$G_{x+y}(z) = \frac{1}{36}(z^2 + 2z^3 + 3z^4 + 4z^5 + 5z^6 + 6z^7 + 5z^8 + 4z^9 + 3z^{10} + 2z^{11} + z^{12})$$

Now from the definition of the PGF:  $w = x + y$  ;  $G_w(z) = \sum_{w=0}^{\infty} p(w)z^w$

Hence:

$w$	0	1	2	3	4	5	6	7	8	9	10	11	12
$p(w)$	0	0	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

$$\mu = \sum_{w=0}^{12} wp(w) = \frac{1}{36}(2 + 6 + 12 + 20 + 30 + 42 + 40 + 36 + 30 + 22 + 12)$$

$$\mu = \frac{252}{36} = 7$$

$$\sigma^2 = \sum_{w=0}^{12} w^2 p(w) - \mu^2$$

$$\sigma^2 = \frac{1}{36}(4 + 18 + 48 + 100 + 180 + 294 + 320 + 324 + 300 + 242 + 144)$$

$$\sigma^2 = \frac{1974}{36} - 7^2 = 5\frac{5}{6}$$

This result is consistent with what could be obtained using an *addition table*.

However, the PGF approach is *powerful* since it could be extended to *more than two dice*, or *dice with different numbers etc.*

+	1	2	3	4	5	6	Outcome	Freq	Probability
1	2	3	4	5	6	7	2	1	$1/36 = 0.028$
2	3	4	5	6	7	8	3	2	$2/36 = 0.056$
3	4	5	6	7	8	9	4	3	$3/36 = 0.083$
4	5	6	7	8	9	10	5	4	$4/36 = 0.111$
5	6	7	8	9	10	11	6	5	$5/36 = 0.139$
6	7	8	9	10	11	12	7	6	$6/36 = 0.167$
7	8	9	10	11	12		8	5	$5/36 = 0.139$
8	9	10	11	12			9	4	$4/36 = 0.111$
9	10	11	12				10	3	$3/36 = 0.083$
10	11	12					11	2	$2/36 = 0.056$
11	12						12	1	$1/36 = 0.028$